

Aortic atresia and interrupted aortic arch communicating through external carotid anastomosis

Image

Cite this article: Dangouloff-Ros V, Boddaert N, and Raimondi F (2019) Aortic atresia and interrupted aortic arch communicating through external carotid anastomosis. *Cardiology in the Young* 29: 699–700. doi: [10.1017/S1047951119000908](https://doi.org/10.1017/S1047951119000908)

Received: 21 January 2019

Revised: 25 February 2019

Accepted: 26 March 2019

First published online: 24 May 2019

Keywords:

Aorta; heart defects; congenital; carotid arteries; CT

Author for correspondence:

Dr Volodia Dangouloff-Ros,
Assistance-Publique Hôpitaux de Paris, Hôpital Necker-Enfants Malades, Department of Pediatric Radiology, 149 rue de Sèvres, 75015 Paris, France. Tel: +33-1-44496719; Fax: +33-1-44495170; E-mail: volodia.dangouloff-ros@aphp.fr

Volodia Dangouloff-Ros^{1,2,3,4} , Nathalie Boddaert^{1,2,3,4} and Francesca Raimondi^{1,5}

¹Department of Pediatric Radiology, Hôpital Universitaire Necker Enfants Malades, AP-HP, 149 rue de Sèvres, 75105 Paris, France; ²INSERM U1000, 149 rue de Sèvres, 75015 Paris, France; ³University René Descartes, PRES Sorbonne Paris Cité, 12 rue de l'Ecole de Médecine, Paris, France; ⁴UMR 1163, Institut Imagine, 24 boulevard du Montparnasse, 75015 Paris, France and ⁵Unité Médico-Chirurgicale de Cardiologie Congénitale et Pédiatrique, Centre de référence Malformations Cardiaques Congénitales Complexes – M3C Hôpital Universitaire Necker Enfants Malades, AP-HP, 149 rue de Sèvres, 75105 Paris, France

Abstract

We describe the case of a newborn girl who displayed association of aortic atresia and interrupted aortic arch, with retrograde flow in ascending aorta, through extracranial anastomoses between vertebral arteries (arisen from descending aorta) and external carotids.

Association of aortic atresia and interrupted aortic arch is usually lethal due to coronary hypoperfusion, unless coronary flow is provided by anastomosis with pulmonary vessels^{1–4}. A girl had prenatal diagnosis of conoventricular ventricular septal defect, aortic atresia, and probable aortic arch interruption. Neonatal echocardiography confirmed prenatal diagnosis, with large ductus arteriosus vascularising descending aorta. During the first day of life, she presented myocardial ischemia, with troponin elevation at 3884 ng/L.

Cervico-thoracic angio-CT scanner displayed aortic valve atresia with hypoplastic ascending aorta connected to common carotid arteries. Aortic arch was interrupted after carotid arteries. Left-sided descending aorta was connected to a large pulmonary trunk through a large ductus arteriosus. Right and left subclavian arteries arise from descending aorta, right subclavian artery having retro-esophageal course, without Kommerell diverticulum. Large right and left vertebral arteries arise from subclavian arteries. On both sides, they vascularised large collateral arteries arising from the junction of their V3 and V4 segments, which were connected to the external carotid arteries. Therefore, coronary arteries were supplied by retrograde flow in common carotids arteries coming from subclavian arteries through an anastomosis of vertebral arteries and external carotid arteries (Fig 1).

At 7 days of age, the patient underwent surgical repair, with pulmonary homograft and Damus–Kaye–Stansel procedure to reconstruct aortic arch, surgical connection of right ventricle to pulmonary arteries using a prosthetic conduit.





Figure 1. 3D reconstruction of arterial system. Anterior (left image), right (middle image), and posterior (right image) view of aortic, supra-aortic, and pulmonary arteries. In red, hypoplastic ascending aorta is connected to common carotid arteries, which terminates as external carotid arteries without internal carotids arteries. In blue, pulmonary artery is connected with descending aorta through a large patent ductus arteriosus; descending aorta is connected with subclavian arteries which continue as two large vertebral arteries. Anastomoses between these two systems are located on both sides at the junction of V3 and V4 segments of vertebral arteries, allowing a retrograde flow within carotid arteries and ascending aorta.

We report here the first case of bilateral extracranial anastomosis between distal vertebral arteries and external carotid arteries, providing retrograde flow to carotids arteries, ascending aorta, and finally to coronary arteries.

Author ORCIDs. Volodia Dangouloff-Ros  [0000-0003-1260-5886](https://orcid.org/0000-0003-1260-5886)

Acknowledgements. None.

Financial Support. This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

Conflict of Interest. None.

Ethical Standards. The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national guidelines on

human experimentation and with the Helsinki Declaration of 1975, as revised in 2008, and has been approved by the institutional committees.

References

1. Careddu L, Oppido G, Ialonardi M, Gargiulo G. Aortic atresia with interrupted aortic arch: a combination incompatible with life? *Interact Cardiovasc Thorac Surg* 2015; 21: 272–273.
2. Tannous HJ, Moulick AN, Jonas RA. Interrupted aortic arch and aortic atresia with circle of Willis-dependent coronary perfusion. *Ann Thorac Surg* 2006; 82: e11–e13.
3. Maruo A, Oshima Y, Hasegawa T. Left subclavian artery perfusion during the Norwood operation for interrupted aortic arch with aortic atresia. *Ann Thorac Surg* 2016; 101: 764–766.
4. Yew G, Coleman D, Calder L. Aortic valvar atresia, interrupted aortic arch, and quadricuspid pulmonary valve: a rare combination. *Pediatr Cardiol* 2005; 26: 455–459.