




I'm not robot



Continue

## Chipmunk cheeks cartoon

The pills with U27 imprint are Blue, Round and have been identified as Amphetamine and Dextroamphetamine 10 mg. This is provided by Aurolife Pharma, LLC. Amphetamines/dextroamphetamine are used in the treatment of adhd; narkolehi and belongs to the cns class of stimulant drugs. Risks cannot be ruled out during pregnancy. Amphetamine/dextroamphetamine 10 mg is classified as Schedule 2 controlled substances under the Controlled Substance Act (CSA). Images for U27 Related Images for U27 More informationAlways consult with your healthcare provider to make sure the information displayed on this page applies to your personal circumstances. Medical Disclaimer dextroamphetamine-amphetamine 5 mg tablet dextroamphetamine-amphetamine 5 mg tablet color dark blue shape round imprint logo and 84 dextroamphetamine-amphetamine 30 mg tablet dextroamphetamine-amphetamine 30 mg tablet color dark orange shape round imprint logo and 404 dextroamphetamine-amphetamine 10 mg tablet dextroamphetamine-amphetamine 10 mg tablet color dark blue shape round imprint logo and 111 dextroamphetamine-amphetamine 20 mg tablet dextroamphetamine-amphetamine 20 mg tablet color dark orange shape round imprint logo and 401 dextroamphetamine-amphetamine 5 mg tablet dextroamphetamine-amphetamine 5 mg tablet color light blue shape round imprint M A5 dextroamphetamine-amphetamine 10 mg tablet dextroamphetamine-amphetamine 10 mg tablet color dark blue shape round imprint M A10 dextroamphetamine-amphetamine 20 mg tablet dextroamphetamine-amphetamine 20 mg tablet color peach shape round imprint M A20 dextroamphetamine-amphetamine 7.5 mg tablet dextroamphetamine-amphetamine 7.5 mg tablet color blue shape round imprint M A7 dextroamphetamine-amphetamine 15 mg tablet dextroamphetamine-amphetamine 15 mg tablet color peach shape round imprint M A30 dextroamphetamine-amphetamine 15 mg tablet color peach shape round imprint M A30 dextroamphetamine-amphetamine 12.5 mg tablet dextroamphetamine-amphetamine 12.5 mg tablet color peach shape oval imprint b 775, 7 1/2 dextroamphetamine-amphetamine 15 mg tablet dextroamphetamine-amphetamine 15 mg tablet color peach shape round imprint b 974, 3 0 dextroamphetamine-amphetamine 5 mg tablet dextroamphetamine-amphetamine 5 mg tablet color blue shape oval imprint b 973, 2 0 15 mg tablet tablet 15 mg tablets bright orange form round imprint U29 dextroamphetamine-amphetamine 20 mg tablets dextroamphetamine-amphetamine 20 mg tablets light orange color imprint U30 dextroamphetamine-amphetamine 5 mg tablets dextroamphetamine-amphetamine 5 mg light blue tablet color round form print U25 dextroamphetamine-amphetamine 10 mg tablets dextroamphetamine-amphetamine 10 mg tablets light blue form round trace U27 dextroamphetamine-amphetamines 30 mg tablets dextroamphetamine-amphetamine 30 mg tablets light orange color round form imprint U31 dextroamphetamine-amphetamine 20 mg tablets dextroamphetamine-amphetamine 20 mg tablets yellow round form imprint MP 446 dextroamphetamine-amphetamine 10 mg dextroamph tabletsetamine-amphetamine 10 mg green color round form imprint MP 443 dextroamphetamine-amphetamine 30 mg tablets dextroamphetamine-amphetamine 30 mg tablets yellow round form imprint MP 4 47 dextroamphetamine-amphetamine 12.5 mg tablets dextroamphetamine-amphetamine 12.5 mg peach color tablets round form imprint N32 dextroamphetamine-amphetamine 7.5 mg peach color tablets round form imprint N32 dextroamphetamine-amphetamine 7.5 mg tablets blue round form prints N30 dextroamphetamine-amphetamine 15 mg tablets dextroamphetamine-amphetamine 15 mg tablets color peach shape round imprint N33 dextroamphetamine-amphetamine 20 mg tablets dextroamphetamine-amphetamine 20 mg tablets white octagonal form imprint M, 30 dextroamphetamine-amphetamine 10 mg tablets dextroamphetamine-amphetamine 10 mg tablets white form pillow imprint M, 10 dextroamphetamine-amphetamine 15 mg tablets white octagonal form imprint M, 15 dextroamphetamine-amphetamine 30 mg tablets dextroamphetamine-amphetamine 30 mg tablets orange color form , 8 0 dextroamphetamine-amphetamine 10 mg tablets dextroamphetamine-amphetamine 10 mg tablets blue round form imprint A, 7 6 dextroamphetamine-amphetamine 20 mg tablets dextroamphetamine-amphetamine 20 mg tablets orange red form 7 9 dextroamphetamine-amphetamine 7.5 mg tablets dextroamphetamine-amphetamine 7.5 mg tablets blue imprint oval cast 131 dextroamphetamine-amphetamine 12.5 mg tablets dextroamphetamine--amphetamine 12.5 mg tablets purple round form imprint cast 133 dextroamphetamine-amphetamine 15 mg tablets dextroamphetamine-amphetamine 15 mg tablets pink oval form imprint cast 134 dextroamphetamine-amphetamine 30 mg tablets dextroamphetamine-amphetamine 30 mg tablets pink round form imprint logo and 345 dextroamphetamine-amphetamine 5 mg tablets dextroamphetamine-amphetamine 5 mg tablets blue round form imprint logo and 339 15 mg tablet color pink shape oval imprint logo and 343 dextroamphetamine-amphetamine 7.5 mg tablet dextroamphetamine-amphetamine 7.5 mg tablet color blue shape round imprint N 951 dextroamphetamine-amphetamine 10 mg tablet dextroamphetamine-amphetamine 10 mg tablet color blue shape oval imprint N 952 dextroamphetamine-amphetamine 15 mg tablet color peach shape round imprint N 954 dextroamphetamine-amphetamine 5 mg tablet dextroamphetamine-amphetamine 5 mg tablet color blue shape oval imprint N 950 dextroamphetamine-amphetamine 12.5 mg tablet color peach shape oval imprint N 953 dextroamphetamine-amphetamine 20 mg tablet dextroamphetamine-amphetamine 20 mg tablet color peach shape oval imprint N 955 dextroamphetamine-amphetamine 30 mg tablet dextroamphetamine-amphetamine 30 mg tablet color peach shape oval imprint N 956 dextroamphetamine-amphetamine 20 mg tablet dextroamphetamine-amphetamine 20 mg tablet color pink shape round imprint logo and 341 dextroamphetamine-amphetamine 30 mg tablet dextroamphetamine-amphetamine 30 mg tablet color pink shape round imprint cast 136 dextroamphetamine-amphetamine 20 mg tablets dextroamphetamine-amphetamine 20 mg pink tablets round form imprint cast 135 dextroamphetamine-amphetamine 10 mg dextro tabletsamphetamine-amphetamine 10 mg tablets blue round form imprint cast 130 See also How to Use parts. Drug interactions can change the way your medication works or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonrepresentative medications and herbal products) and share them with your doctor and pharmacist. Do not initiate, discontinue, or change the dosage of any medication without your doctor's approval. The drug can interact with many products. Some products include: apraclonidine, atomoxetine, bupropion, carbamazepine, dichlorphenamide, methotrexate, mifepristone, potassium capsules / tablets, pramlintide, tetrabenazine, certain Parkinson's disease drugs (entacapone, levodopa, tolcapone), certain high blood pressure drugs (guanethidine, methyldopa), sulfonamide drugs (including sulfa antibiotics such as sulfamethoxazole), sympathizers (such as amphetamines, ephedrine), products that reduce the amount of acid in the urine (such as antacids, sodium bicarbonate, potassium / sodium citrate, acetazo This drug can increase the risk of bleeding when processed with other drugs that can also cause bleeding. Examples include anti-platelet drugs such as clopidogrel, thinners such as dabigatran / exoxaparin / warfarin, among others. Taking other MAO inhibitors with these drugs can lead to serious (possibly fatal) drug interactions. Avoid taking other MAO inhibitors (isocarboxazid, linezolid, moclobemide, fenelzine, procarbazine, rasagiline, selegiline, tranlycypromine) during treatment with this drug. Most MAO inhibitors should also not be minimized for two weeks before and after treatment with this drug. Ask your doctor when to start or stop taking this medication. Before using this product, report the use of other medications that can increase the risk of very high blood pressure (hypertensive crisis) when combined with this product, including herbal products (such as ephedra /ma huang), nasal decongestants (such as fenefesfrin, pseudoephedrine), stimulants (such as amphetamines, ephedrine, epinephrine), dietary aids, among others. This product should not be used with any of these drugs. Talk to your doctor or pharmacist for more details. If you also take antifungal drugs azole (such as itraconazole, ketoconazole), take this product at least 2 hours after the antifungal drug. Check all prescription and nonprescription drug labels carefully as many medications contain pain reliever/fever reliever (aspirin, NSAIDs such as ibuprofen, ketorolac or naproxen). These drugs are similar to salicylate and can increase your risk of side effects if taken together. However, if your doctor has directed you to take low-dose aspirin to prevent a heart attack or stroke (usually at a dose of 81-325 milligrams a day), you should continue taking aspirin unless your doctor instructs you otherwise. Ask your doctor or pharmacist for more details. Tell your doctor or pharmacist if you take other products that cause drowsiness including alcohol, marijuana (marijuana), antihistamines (such as cetirizine, diphenhydramine), medications for sleep or anxiety (such as alprazolam, diazepam, zolpidem), muscle relaxants, and opioid pain reliever (such as codeine). Check the labels on all your medications (such as allergies or cough and cold products) as they may contain ingredients that cause drowsiness. Ask your pharmacist about using these products safely. The drug can interfere with certain laboratory tests, possibly leading to false test results. Make sure the laboratory personnel and all your doctors know you are using this drug. Page 2 Before taking quetiapine, tell your doctor or pharmacist if you are allergic to it; or if you have other allergies. The product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using the drug tell your doctor or pharmacist your medical history, especially about: eye cataracts, liver disease, low white blood cell count (including a history of low white blood cell count caused by seizure disorders, difficulty swallowing, thyroid problems, gastric / intestinal blockages (such as severe constipation, intestinal obstruction), stomach/intestinal motion (such as ileus), personal or family history of diabetes, personal or family history of substance use disorders (such as excessive use of drug/alcohol addiction), heart disease, high blood pressure, high cholesterol/triglyceride levels, sleep apnea problems, difficulty urinating (for example, due to enlarged prostate). The drug can make you dizzy or sleepy or obscure your vision. Alcohol or marijuana (marijuana) can make you more dizzy or sleepy. Do not drive, use the engine, or do anything that requires vigilance or clear vision until you can do so safely. Avoid alcoholic beverages. Talk to your doctor if you use marijuana. Quetiapine can cause conditions that affect heart rhythm (QT extension). The extension of QT can rarely cause serious (rarely fatal) rapid/irregular heartbeats and other symptoms (such as severe dizziness, fainting) that require immediate medical attention. The risk of QT extension may increase if you have certain medical conditions or take other medications that may cause QT renewal. Before using quetiapine, tell your doctor or pharmacist about all the medications you take and if you have any of the following conditions: certain heart problems (heart failure, slow heart rate, QT extension in ECG), family history of certain heart problems (QT extension in ECG, sudden cardiac death). Low levels of potassium or magnesium in the blood can also increase your risk of prolonging QT. This risk may increase if you use certain medications (such as diuretics/water pills) or if you have conditions such as severe sweating, diarrhea, or vomiting. Talk to your doctor about using quetiapine safely. This drug may make you sweat less, making you more likely to have a heat stroke. Avoid doing things that can cause you to overheat, such as working hard or exercising in hot weather, or using a hot tub. When the weather is hot, drink plenty of fluids and dress lightly. If you are too hot, quickly find a place to cool down and rest. Get immediate medical help if your fever is not gone, mental/mood swings, headaches, or dizziness. Before undergoing surgery, tell your doctor or dentist about all the products you use (including prescription medications, non-incription medications, and herbal products). Children may be at higher risk for certain side effects when taking these medications, such as increased blood pressure or prolactin (see also Side Effects section). Talk to your doctor about the risks and benefits. Older adults may be more sensitive to side effects of this drug, especially drowsiness, dizziness, lightheadness, and the extension of QT (see above). Drowsiness, dizziness, and lightheadness can increase the risk of pregnancy, the drug should be used only when it is clearly needed. Babies born to mothers who used this drug during the last 3 months of pregnancy may rarely experience symptoms including muscle stiffness or shaking, drowsiness, difficulty eating/breathing, or crying constantly. If you notice any of these symptoms in your newborn especially during their first month, immediately notify the doctor. Since untreated mental/mood problems (such as schizophrenia, bipolar disorder, depression) can be a serious condition, do not stop taking these medications unless directed by your doctor. If you are planning a pregnancy, getting pregnant, or thinking you may be pregnant, immediately discuss with your doctor the benefits and risks of using this medication during pregnancy. The drug enters breast milk and may have an undesirable effect on nursing babies. Immediately tell your doctor if your baby is experiencing symptoms such as muscle stiffness or shaking, unusual drowsiness, or difficulty feeding. Consult your doctor before breastfeeding. Page 3 See also The Warnings section. Before taking sotalol, tell your doctor or pharmacist if you are allergic to it; or if you have other allergies. The product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this drug, tell your doctor or pharmacist your medical history, especially from: certain heart rhythm problems (such as a slow heartbeat, Sore sinus syndrome, second or third degree atrioventricular block unless you have a pacemaker), kidney problems, heart failure, respiratory problems (such as asthma, chronic bronchitis, emphysema), very recent heart attack (within 2 weeks), overactive thyroid disease (hyperthyroidism), serious allergic reactions requiring treatment with sotalol epinephoid can cause conditions that affect heart rhythm (extension of QT in ECG). The extension of QT can rarely cause serious (rarely fatal) rapid/irregular heartbeats and other symptoms (such as severe dizziness, fainting) that require immediate medical attention. The risk of QT extension may increase if you have certain medical conditions or take other medications that may cause QT renewal. Before using sotalol, tell your doctor or pharmacist about all the medications you take and if you have any of the following conditions: certain heart problems (QT extension in ECG, history of torsade de pointes), family history of certain heart problems (QT extension in ECG, sudden cardiac death). Low levels of potassium or magnesium in the blood can also increase your risk of prolonging QT. This risk may increase if you use certain medications (such as diuretic water pills/water), if you are unable to eat or drink liquids as usual, or if you have conditions such as severe/prolonged sweating, diarrhea, or vomiting. Talk to you to you about using sotalol safely. If you have diabetes, this product can mask the fast heart rate/pounding that you usually feel when your blood sugar falls too low (hypoglycemia). Other symptoms of low blood sugar, such as dizziness and sweating, are not affected by this drug. This product can also make it more difficult to control your blood sugar. Check your blood sugar regularly as directed and share the results with your doctor. Immediately tell your doctor if you have symptoms of high blood sugar such as increased thirst/urination. Your doctor may need to adjust your diabetes medication, exercise program, or diet. Before undergoing surgery, tell your doctor or dentist that you are taking this medication. This drug might make you dizzy. Alcohol or marijuana (marijuana) can make you more dizzy. Don't drive, use the engine, or do anything that requires vigilance until you can do it safely. Limit alcoholic beverages. Talk to your doctor if you use marijuana. Older adults may be more sensitive to side effects of this drug, especially dizziness, fatigue, and QT extension (see above). During pregnancy, the drug should be used only when it is clearly needed. This can harm the unborn baby. Discuss the risks and benefits with your doctor. The drug enters breast milk and may have an undesirable effect on nursing babies. Discuss the risks and benefits with your doctor before breastfeeding. Page 4See also The Warnings section. Drug interactions can change the way your medication works or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonrepresentative medications and herbal products) and share them with your doctor and pharmacist. Do not initiate, discontinue, or change the dosage of any medication without your doctor's approval. Some products that may interact with this drug include: kava, sodium oxidate. Other drugs can affect the removal of alprazolam from your body, which can affect the way alprazolam works. Examples include antifungal azole (such as itraconazole, ketoconazole), cimetidine, certain anti-depressants (such as fluoxetine, fluvoxamine, nefazodone), drugs to treat HIV (delavirdine, protease inhibitors such as indinavir), macrolide antibiotics (such as erythromycin), rifamycins (such as rifabutin), St. John's wort, drugs used to treat seizures (such as erythronin), among others. The risk of serious side effects (such as slow/shallow breathing, severe drowsiness/dizziness) may increase if the drug is taken with other products that can also cause drowsiness or respiratory problems. Tell your doctor or if you take other products such as opioid pain or cough reliever (such as codeine, hydrocodone), alcohol, marijuana (marijuana), other medications for sleep or anxiety (such as diazepam, lorazepam, zolpidem), muscle relaxants as carisoprodol, cyclobenzaprine), or antihistamines (such as cetirizine, diphenhydramine). Check the labels on all your medications (such as allergies or cough and cold products) as they may contain ingredients that cause drowsiness. Ask your pharmacist about using these products safely. Smoking reduces the blood levels of this drug. Let your doctor know if you smoke or if you have recently quit smoking. Page 5 alprazolam ER 1 mg tablets, extended release 24 hours alprazolam ER 1 mg tablets, extended release 24 hours yellow color round shape print R, 84 alprazolam ER 2 mg tablets, extended release 24 hr alprazolam ER 2 mg tablets, extended release 24 hr color peach shape round imprint R, 87 alprazolam ER 0.5 mg tablets, extended release 24 hr alprazolam ER 0.5 mg tablets, extended release 24 hr white round shape imprint logo, 83 alprazolam ER 3 mg tablets, extended release 24 hours alprazolam ER 3 mg tablets, extended release 24 hours light green color round shape imprint R, 86 alprazolam ER 0.5 mg tablets, extended release 24 hours alprazolam ER 0.5 mg tablets, extended release 24 hr white round shape imprint cast 187 alprazolam ER 2 mg tablets, extended release 24 hr alprazolam ER 2 mg tablets, extended release 24 hr blue round shape G print, 2 alprazolam ER 1 mg tablets, extended release 24 hr alprazolam ER 1 mg tablets, extended release 24 hr yellow color imprint IP 10 alprazolam ER 1 mg tablets, extended release 24 hr alprazolam ER 1 mg tablets, extended release 24 hr yellow square shape . 1 alprazolam ER 2 mg tablets, extended release 24 hours alprazolam ER 2 mg tablets, extended release 24 hr blue oval shape imprint 93, 5452 alprazolam ER 3 mg tablets, extended release 24 hr alprazolam ER 3 mg tablets, extended release 24 hr green round shape imprint 93, 5453 alprazolam ER 1 mg tablets , extended release 24 hr yellow round shape imprint 93, 5451 alprazolam ER 0.5 mg tablets, extended release 24 hours white round shape imprint 93, 5450 alprazolam ER 1 mg tablets, extended release 24 hr yellow round shape imprint cast 188 alprazolam ER 0.5 mg tablets, extended release 24 hr alprazolam ER 0.5 mg tablets, extended release 24 hr white round shape X print, 70 alprazolam ER 1 mg tablets, extended release 24 hr alprazolam ER 1 mg tablets,extended release 24 hr yellow form round imprint x, 73 alprazolam ER 2 mg tablets,extended release 24 hr blue round shape imprint Xprint X, 74 alprazolam ER 3 mg tablets, extended release 24 hr alprazolam ER 3 mg tablets,extended release 24 hr green color round shape imprint X, 75 alprazolam ER 3 mg tablets,extended release 24 hr alprazolam ER 3 mg release 24 hr green color Traces of G, 3 alprazolam ER 0.5 mg tablets, extended release 24 hr alprazolam ER 0.5 mg tablets, extended release 24 hr white form pentagon imprint G, 0.5 Page 6 Drugs amp; Drugs Using Side Effects Of Overdose Interaction Action Images Using Side Effects Of Overdose Interaction Precautions Images Using Carisoprodol used short term to treat Usually used along with rest, physical therapy, and other treatments. It works by helping to relax the muscles. Take this medicine by mouth with or without food as directed by your doctor. Do not increase your dosage or use this drug more often or longer than specified. Your condition will not improve faster, and the risk of your side effects will increase. Dosage is based on your medical condition and response to treatment. The drug should only be used short term (for 3 weeks or less) unless directed by your doctor. If you suddenly stop using this medication, you may have withdrawal symptoms (such as abdominal cramps, difficulty sleeping, headache, nausea). To help prevent withdrawal, your doctor can lower your dosage slowly. Withdrawal is more likely if you have been using carisoprodol for a long time or in high doses. Immediately notify your doctor or pharmacist if you have any withdrawals. Although it helps many people, this drug can sometimes cause addiction. This risk may be higher if you have substance use disorders (such as overdose or drug/alcohol addiction). Take this drug exactly as prescribed to lower the risk of addiction. Ask your doctor or pharmacist for more details. Tell your doctor know if your condition persists after 2 to 3 weeks or if it worsens. Side Effects Dizziness, drowsiness, or headaches may occur. If any of these effects persist or worsen, immediately notify your doctor or pharmacist. Remember that your doctor has prescribed this medication because he has judged that the benefits for you outweigh the risk of side effects. Many people who use this drug do not have serious side effects. Immediately notify your doctor if you have serious side effects, including: confusion. A very serious allergic reaction to the drug is rare. However, get medical help immediately if you notice symptoms of a serious allergic reaction, including: rash, itching / swelling (especially the face / tongue / throat), severe dizziness, difficulty breathing. This is not a complete list of possible side effects. If you notice any other effects not listed above, contact your doctor or pharmacist. In the US -Contact your doctor for medical advice on side effects. You can report side effects to the FDA at or in www.fda.gov/medwatch.In Canada – Contact your doctor for medical advice on side effects. You can report any side effects to Health Canada at 1-866-234-2345. Precautions Before taking carisoprodol, tell your doctor or pharmacist if you are allergic to it; or to tybamate, or blinding; or if you have other allergies. The product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this drug, tell your doctor or pharmacist your medical history, especially about: certain blood disorders (acute intermittent porphyria), kidney disease, liver disease, seizures, personal or family history of substance use disorders (such as overdose or drug/alcohol addiction). This remedy may make you dizzy or sleepy. Alcohol or marijuana (marijuana) can make you more dizzy or sleepy. Don't drive, use the engine, or do anything that requires vigilance until you can do it safely. Avoid alcoholic beverages. Talk to your doctor if you use marijuana. Before undergoing surgery, tell your doctor or dentist about all the products you use (including prescription medications, non-incription medications, and herbal products). Older adults may be more sensitive to side effects of this drug, especially drowsiness, or confusion. These side effects can increase the risk of falls. During pregnancy, the drug should be used only when it is clearly needed. Discuss the risks and benefits with your doctor. The drug enters breast milk and may have an undesirable effect on nursing babies. Consult your doctor before breastfeeding. Drug Interaction interactions may change the way your medication works or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonrepresentative medications and herbal products) and share them with your doctor and pharmacist. Do not initiate, discontinue, or change the dosage of any medication without your doctor's approval. Tell your doctor or pharmacist if you are taking other products that cause drowsiness such as opioid pain or cough reliever (such as codeine, hydrocodone), alcohol, marijuana (marijuana), medications for sleep or anxiety (such as alprazolam, lorazepam, zolpidem), other muscle relaxants (such as cyclobenzaprine, methocarbamol), or antihistamines (such as cetirizine, dithydramine). Check the labels on all your medications (such as allergies or cold products) as they may contain ingredients that cause drowsiness. Ask your pharmacist about using these products safely. Overdose If someone has overdosed and has serious symptoms such as fainting or difficulty breathing, call 911. If not, contact the poison control center immediately. U.S. residents can call their local poison control center at 1-800-222-1222. Canadians can call the province's poison control center. Symptoms of an overdose can be severe drowsiness/dizziness, seizures, slow/shallow breathing, mental/mood swings (such as confusion, hallucinations), inability to move your legs/arms, wobbly/wobbly movements, vision changes (such as visions). NoteA drug is not shared with others. Sharing is against the law. The drug has been prescribed for your current condition only. Don't use it later for other conditions unless your doctor directs you to do so. Different drugs may be necessary in that case. Dose missedIf you miss the dosage, take it as soon as possible. If it is nearing the next dose time, skip the missed dose. Take the next dose at regular times. Don't double the dosage to catch up. StorageStore at room temperature is far from light and humidity. Do not store in the bathroom. Keep all drugs from children and pets. Do not flush the drug into the toilet or pour it into the drain unless instructed to do so. Dispose of this product correctly when it has expired or is no longer needed. Consult your local pharmacist or waste disposal company.The information was last revised april 2019. Copyright(c) 2019 First Databank, Inc. Image carisoprodol 350 mg tablet carisoprodol 350 mg tablets white round shape imprint 2410 V carisoprodol 350 mg tablets carisoprodol 350 mg tablets white round form imprint SG 109 carisoprodol 350 mg tablets carisoprodol 350 mg tablets white round form imprint 5513, AND carisoprodol 350 mg tablets carisoprodol 350 mg tablets white round form imprint CL 022 carisoprodol 350 mg tablets carisoprodol 350 mg tablets white round shape 111 . logo carisoprodol 350 mg tablets carisoprodol 350 mg tablets white round form imprint 446 carisoprodol 350 mg tablets carisoprodol 350 mg tablets white round form imprint OP 35 carisoprodol 250 mg tablets carisoprodol 2250 mg white tablets round form imprint S 434 carisoprodol 250 mg tablets carisoprodol 250 mg tablets white round form imprint WP 5901 carisoprodol 350 mg tablet carisoprodol 350 mg tablets white round form imprint D. 31 carisoprodol 350 mg tablet carisoprodol 350 mg tablets color white shape round imprint WW 176 More Resources for Carisoprodol Find Lowest Prices on Check Interactions Read Reviews (239) Selected from data included with permission and copyrighted by First Databank, Inc. This copyrighted material has been downloaded from licensed data providers and is not for distribution, except as permitted by applicable terms of use. CONDITIONS OF USE: The information in this database is intended to complement, not replace, the expertise and assessment of health professionals. This information is not intended to cover all possible uses, referrals, precautions, drug interactions or side effects, nor should it be construed to indicate that the use of a particular drug is safe, appropriate or effective for you or others. A health care professional should before taking any medication, change any diet or start or discontinue any treatment. Treatment.