




I'm not robot



Continue

Progon m capsule side effects

Manufacturer/Marketer Address21/2,Mc Nichols Road, Chetpet, Chennai-600 031, India Country of Origin: India Indication Men's and Women's Infertility. Dosage1 capsule three times a day or according to your doctor's instructions. How long to spend? This product can be used for 2-3 months on the advice of your doctor. With Western medicines Ask your doctor for advice if you are taking this product in combination with other Western (allopathic/ modern) medicines. Some ayurveda herbs can interact with modern medicine. If both ayurveda and allopathic medicines are recommended together, it is best to take an allopathic medicine first, wait 30 minutes and then, after a gap of 15 to 30 minutes, take ayurveda medicine Can this be used during use of homeopathic medicine? Yes. This product does not react with homeopathic medicine. With supplements such as multivitamin tablets, Omega 3 fatty acids, etc?? Yes. In general, this product is well suited with most food supplements. However, if you take more than one product a day, contact your doctor. IngredientsEach capsule contains – Ashwagandha – Indian ginseng – Withania somnifera – Aphrodisiac, nervine tonic, corrects male seminal instability & impotence. Yastimadhu – Licorice – Glycyrrhiza glabra – Improves libido, increases sperm volume, corrects fatigue. Jatiphala – Nutmeg – Myristica Fragrances – Improves libido, corrects premature ejaculation and erection. Gokshura – Tribulus terrestris – Restorative tonic for vitality and vitality. Payaswini – Vidarikanda – Ipomea digitata – Anxilytic, improves psychological trust. Musali – Curculigo orchioides – An excellent rejuvenation, drains the seed channels. Somavalli – Guduchi – Giloy – Tinospora cordifolia – Addresses idiopathic sexual inadequacy and sterility. Side effects progon M Capsule This medicine has no known side effects. However, it is best to use this drug under the supervision of a doctor. Store in a cool dry place, away from sunlight. Stay out of the reach and sight of children. Packaging, manufacturer, shelf lifePresentation & packaging:10 capsule blister. 30 capsule bottle. Manufacturer: Retort Pharma Pvt Ltd.Shelf life: 3 years from the date of manufacture. Generic name: Progesterone Capsules (proe JES ter one)Brand name: Prometrium Medically reviewed by Drugs.com. Last updated: 7.6.2020. Warning Do not use estrogen with progestin like this medicine (progesterone capsules) to prevent heart disease or dementia. The use of estrogens with progestin may increase the chance of heart attack, stroke, breast cancer, blood clots or dementia. Use estrogen with or without progestin for the shortest possible time at the lowest dose. It's used. the possibility of changes in the lining of the uterus in women life change receiving estrogen therapy. It is used to treat women without a monthly cycle. It can be given to you for other reasons. Talk to the doctor. If you have an allergy to progesterone or any of the other ingredients of this medicine (progesterone capsules). If you have a peanut allergy and take the capsules. There's peanut oil in the capsules. If you are allergic to this medicine (progesterone capsules); any part of this drug (progesterone capsules); or other medicines, food or substances. Tell your doctor about the allergy and what signs you had. If you have had any of these health problems: Bleeding disorder; the risk of blood clots or blood clots; breast cancer or other cancer; liver disease; heart attack; stroke; or a tumor in which estrogen or progesterone causes it to grow. If you have unexplained vaginal bleeding. If you're pregnant or you might be pregnant. Do not take this medicine (progesterone capsules) if you are pregnant. This is not a list of all drugs or health problems that interact with this drug (progesterone capsules). Tell your doctor and pharmacist about all your medicines (prescription or OTC, natural products, vitamins) and health problems. Check that it is safe for you to take this medicine (progesterone capsules) with all medicines and health problems. Do not start, stop or change the dose without consulting your doctor. What do I need to know or do when I take Progesterone capsules? Tell all healthcare providers that you are taking this medicine (progesterone capsules). This also applies to doctors, nurses, pharmacists and dentists. Avoid driving and other tasks or actions that require you to be vigilant until you see how this drug (progesterone capsules) affects you. If you are allergic to any type of nuts or seeds, talk to your doctor. If you have high blood sugar (diabetes), talk to your doctor. This medicine can raise blood sugar levels. Make sure you have regular breast exams and gynecology checks. You also need to do breast self-examinations, as you have been told. This drug can affect certain laboratory tests. Tell all healthcare providers and laboratory workers that you are taking this medicine (progesterone capsules). A blood clot has occurred with this medicine (progesterone capsules). These clots have included a heart attack, stroke and clot in the leg, lungs or eye. Sometimes a blood clot can be deadly. Tell your doctor if you have ever had a blood clot. Talk to your doctor. Talk to your doctor if you need to spend long periods of time, such as long distances, bedrest after surgery or illness. Not moving for long periods of time can add Possibility. If you are 65 years of age or older, use this medicine (progesterone capsules) carefully. You could have more side effects. This medicine is not approved for use in children. Children. With a doctor. This medicine may prevent menstruation (menstrual bleeding) for some time. This is not a method of contraception. Tell your doctor if you are breast-feeding. You need to talk about all the risks to your baby. How is this medicine (Progesterone Capsules) best used? Use this medicine (progesterone capsules) at the behest of your doctor. Read all the information you've been given. Follow all instructions carefully. Take it at bedtime if you take it once a day. Take with or without food. Take with food if it causes stomach problems. Have a glass of water. If you have difficulty swallowing, talk to your doctor. What do I do if I don't get a dose? Take the missed dose as soon as you think about it. If the time for the next dose is near, skip the missed dose and return to your standard time. Do not take 2 doses at the same time or extra doses. What are the side effects that I need to call my doctor right away? WARNING/WARNING: Although it may be rare, some people may have very bad and sometimes deadly side effects while taking the drug. Tell your doctor or get medical advice immediately if you have any of the following symptoms or symptoms that may be associated with a very bad side effect: Signs of an allergic reaction, such as a rash; nettle dosing; itching; red, swollen, blistering or exfoliating skin with or without fever; wheezing; tightness in the chest or throat; difficulty breathing, swallowing or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue or throat. Signs of liver problems such as dark urine, fatigue, no hunger, stomach problems or abdominal pain, light stool, vomiting or yellow skin or eyes. Weakness on one side of the body, problems with talking or thinking, change of balance, drooping on one side of the face or blurred vision. Shortness of breath, high weight gain or swelling in the arms or legs. It's hard to walk. Very bad dizziness or fainting. Vision changes or disappears, bulging eyes or changes near contact lenses. Lump in the chest, chest pain or soreness or a burst of the tin. Vaginal itching or discharge. Vaginal bleeding is not normal. Pain when urinating. Urinating more often. Depression or other mood changes. Memory problems or losses. Scenes. Redness. Fast or abnormal heartbeat. This medicine can cause fluid to swell or keep liquid in your body. Tell your doctor if you have swelling, weight gain or difficulty breathing. Call your doctor immediately if you have any signs of a blood clot, such as chest pain or pressure; coughs up blood; shortness of breath; swelling, heat, numbness, discoloration or pain in the leg or arm; or difficulty talking or swallowing. What are the other side effects with Progesterone Capsules? All medicines can cause side effects. A lot of people don't have no side effects or have only minor side effects. Contact your doctor or get your doctor's help if any of these side effects bother you or do not go away: Dizziness, drowsy, tired or weak. Sleeping. Hungrier. Headache. Pimples (acne). Hair loss. Hair growth. Diarrhea or constipation. Stomach problems or vomiting. Abdominal pain or cramps. Swelling. Enlarged breasts. They're a good breast. Back, muscle or joint pain. Weight gain or weight loss. Vaginal bleeding or spotting. Less interest in sex. These are not all side effects that may occur. If you have any questions about side effects, ask your doctor. Call your doctor to ask your doctor about side effects. You can report adverse reactions to the FDA 1-800-332-1088. You can also report side effects. If an overdose is suspected: If you think an overdose has occurred, call the Anti-Poison Control Center or get medical attention immediately. Be prepared to tell or show what was taken, how much and when it happened. How do I store progesterone capsules and/or throw them away? Store at room temperature. Store in a dry place. Do not store in the bathroom. Keep all the drugs in a safe place. Keep all drugs out of the reach of children and pets. Discontinue unused or expired medicines. Do not flush down the toilet or pour down the drain unless you are told to do so. Check with the pharmacy to see if you have any questions about the best agglomeration to throw drugs away. There may be drug recovery programs in your area. Using consumer information If your symptoms or health problems do not improve or worsen, contact your doctor. Do not share your medication with others or take anyone else's medication. Some medicines may have another medical bulletin. Check with the pharmacy. If you have any questions about this medicine (progesterone capsules), talk to your doctor, nurse, pharmacist or other healthcare provider. If you think there's been an overdose, call the poison center or get medical attention right away. Be prepared to tell or show what was taken, how much and when it happened. For more information, please contact your healthcare provider to ensure that the information displayed on this page applies to your personal circumstances. Medical disclaimer