

44 159 pill side effects Active ingredient(s): Paracetamol, Inactive ingredients of aspirin and caffeine: HYPROMELLOSESCELLULOSE, MICROCRYSTALLINEPROPYLENE GLYCOLSODIUM LAURYL SULFATESTEARIC ACIDTITANIUM DIOXIDESTARCH, CORNCROSPOVIDONE View NDC 57243-159 Product type: Human Otc drug dosage form: tablet, film coated printing: 44;159 What is the footprint? The footprint is a feature of an oral solid dosage form of a drug, specifying the alphanumeric text that appears in the form of solid dosing, including text that is engraved, engraved, engraved, engraved, engraved, engraved, or printed in ink. A semicolon is used to show the separation between words or line divisions. Color(s): ColorThe color is a characteristic of an oral solid dosage form of a drug, specifying the color or colors that most define the appearance of the dosage form. Color is not an FDA specification for the actual color of solid dosage shapes or the names of colors that may appear on the product label. Shape: Round (C48348); ShapeThe shape is a characteristic of an oral solid dosage form of a drug, specifying the two-dimensional representation of the solid dose shape, in terms of the outer perimeter of a solid dosage form when the dosage form when the dosage form, supported by a flat surface, is seen directly from above, including a slight rounding of the corners. The shape does not include relief, punctuation, cuts, or internal clippings. The shape is independent of the print orientation and logo. Size: 11 mm (0.43 in) SizeThe size is a characteristic of an oral solid dosage form as a physical amount in the length dimension. The size is indicated in millimeters and rounded to the nearest whole millimeter. Rating: 1 What is Score? Scoring is the number of pieces that a pill could be divided into using the scoring lines present in the pill. For example: (i) A pill with a single score line could be divided into two parts; therefore, it has a score value of 4. iii) A pill without scoring lines has a score value of 1. CVS.com® is not available to customers or patients outside the United States or U.S. territories We apologize for any inconvenience. For U.S. military personnel permanently assigned or temporary service abroad, call our Customer Service team at 1-800-SHOP CVS (1-800-746-7287) if you need help with your Pill with print 44 159 is white, round and has been identified as acetaminophen, aspirin and caffeine 250 mg / 250 mg / 250 mg / 65 mg. It is supplied by Teva Pharmaceuticals USA. Acetaminophen/aspirin/caffeine is used in the treatment of migraine; osteoarthritis; pain/fever and belongs to the analgesic combinations of the drug class. FDA has not classified the risk during pregnancy. Paracetamol/aspirin/caffeine 250 mg / 250 mg / 250 mg / 65 mg is not a controlled Substance under the Controlled Substances Act (CSA). Images for 44 159 Other Brands: Excedrin, Excedrin, Excedrin Migraine, Backaid Inflammatory Pain Formula, Genace, ... +6 More Professional Resources Treatment Guides Related to Migraine Osteoarthritis Pain/Fever Learn moreif you consult your health care provider to ensure that the information shown on this page applies to your personal circumstances. IMPORTANT Medical Disclaimer: HOW TO USE THIS INFORMATION: This is a summary and does NOT have as much information as possible about this product. This information does not guarantee that this product will be safe, effective or appropriate for you. This information is not an individual medical advice and is not a substitute for your health care professional's advice. Always ask your health needs. An ingredient in this product is acetaminophen. Taking too much acetaminophen can cause severe (possibly fatal) liver disease. Adults should not take more than 4000 milligrams (4 grams) of acetaminophen without first asking your doctor or pharmacist. Acetaminophen is in many over-the-counter and prescription medications (such as pain relievers/fever drugs or cough and cold products). Check the labels for all your medicines to see if they contain acetaminophen and ask your pharmacist if you are not sure. Get medical help right away if you take too much acetaminophen (overdose), even if you feel well. Symptoms of overdose may include nausea, vomiting, loss of appetite, sweating, stomach/abdominal pain, extreme tiredness, yellowing of the eyes/skin and dark urine. Daily alcohol consumption, especially when combined with acetaminophen, can damage the liver. Avoid alcohol. This product is a combination of aspirin, acetaminophen and caffeine. It is used for temporary pain relief from conditions such as muscle aches, toothache, menstrual cramps, or headaches (including migraine). Aspirin and acetaminophen relieve pain by preventing the body from containing certain natural substances. Caffeine helps increase the effects of aspirin and acetaminophen. See also the Warning section. If you are taking the over-the-counter self-treating product, read all instructions on the product package before taking this medicine. If you have any questions, ask your doctor, or follow the instructions on the product package. Drink a full glass of water (8 ounces/240 milliliters) with this medicine unless your doctor tells you to Otherwise. If you are using the tablet form of this medicine, do not lie down for at least 10 minutes after taking this medicine, do not lie down for at least 10 minutes after taking this medicine. If you are using the tablet form of this medicine, do not lie down for at least 10 minutes after taking this medicine. If you are using the tablet form of this medicine. If you are using the tablet form of this medicine, do not lie down for at least 10 minutes after taking this medicine. If you are using the tablet form of this medicine and take it with food or milk. The dose is based on your medical condition and response to treatment. Do not increase your dose or use this medicine more often or for longer than 10 days or a fever that lasts longer than 3 days. Use the smallest effective dose. Talk to your doctor or pharmacist for more information. If you are taking this medicine as needed (not on a regular schedule), remember that pain relievers work best if used when the first signs of pain occur. If you wait until the pain has gotten worse, the medication may not work as well. If you are taking this product for headache, seek immediate attention if you also have: trouble speaking, weakness on one side of the body, sudden changes in vision. Before using this product, consult a doctor or pharmacist if you have: daily headaches; his first headache after age 50; a headache that is more severe or different from your usual headaches; headaches; headaches; headaches; headaches; headaches; headache with persistent/severe vomiting, fever and stiffness in the neck. Talk to your doctor or pharmacist if your migraine is not relieved or if it gets worse after the first dose. If you're using migraine attack medications in 10 or more days each month, medications can make your headaches worse (overuse headache). Do not use medicine more often, if the medicine is not working as well, or if your headaches get worse. See your doctor if your condition persists or gets worse (for example, if you have new or unusual symptoms, redness or swelling of the painful area, pain or fever that doesn't go away or get worse). See also the Warning section. Nausea, vomiting, upset stomach, trouble falling asleep or a feeling of tremor/nerves. If any of these effects persist or worsen, tell your doctor or pharmacist promptly. If your doctor has told you to use this medicine, remember that he or she has judged that the benefit to you is greater than the risk of side effects. Many people who use this medicine and tell your doctor right away if any of these unlikely but serious side effects occur: severe dizziness, fainting, persistent stomach pain or vomiting that looks like coffee pots, black/bloody stools, ringing in the ears, hearing loss, irregular/rapid heartbeat, signs of kidney problems (such as changes in the amount of urine), signs of infection (e.g., persistent sore throat or fever), A very severe allergic reaction to this medicine is unlikely, but look for signs of infection (e.g., persistent sore throat or fever), A very severe allergic reaction to this medicine is unlikely, but look for signs of infection (e.g., persistent sore throat or fever), A very severe allergic reaction to this medicine is unlikely, but look for signs of infection (e.g., persistent sore throat or fever), A very severe allergic reaction to this medicine is unlikely, but look for signs of infection (e.g., persistent sore throat or fever), A very severe allergic reaction to this medicine is unlikely, but look for signs of infection (e.g., persistent sore throat or fever), A very severe allergic reaction to this medicine is unlikely, but look for signs of infection (e.g., persistent sore throat or fever), A very severe allergic reaction to this medicine is unlikely, but look for signs of infection (e.g., persistent sore throat or fever), A very severe allergic reaction to this medicine is unlikely, but look for signs of infection (e.g., persistent sore throat or fever), A very severe allergic reaction to this medicine is unlikely, but look for signs of infection (e.g., persistent sore throat or fever), A very severe allergic reaction to this medicine is unlikely, but look for signs of infection (e.g., persistent sore throat or fever). this medicine is unlikely, but look for signs of infection (e.g., persistent sore throat or fever), A very severe allergic reaction to this medicine is unlikely, but look for signs of infection (e.g., persistent sore throat or fever), A very severe allergic reaction to this medicine is unlikely, but look for signs of infection (e.g., persistent sore throat or fever), A very severe allergic reaction to this medicine is unlikely, but look for signs of infection (e.g., persistent sore throat or fever), A very severe allergic reaction to this medicine is unlikely, but look for signs of infection (e.g., persistent sore throat or fever), A very severe allergic reaction to this medicine is unlikely, but look for signs of infection (e.g., persistent sore throat or fever), A very severe allergic reaction to this medicine is unlikely, but look for signs of infection (e.g., persistent sore throat or fever), A very severe allergic reaction to this medicine is unlikely, but look for signs of infection (e.g., persistent sore throat or fever), A very severe allergic reaction to this medicine is unlikely, but look for signs of infection (e.g., persistent sore throat or fever), A very severe allergic reaction to this medicine is unlikely, but look for signs of infection (e.g., persistent sore throat or fever), A very severe allergic reaction to this medicine is unlikely, but look for signs of infection (e.g., persistent sore throat or fever). allergic reaction to this medicine is unlikely, but look for if it happens. Symptoms of a severe allergic reaction may include: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, shortness of breath. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the United States - Call your doctor for medical advice on side effects. You can report side effects to the FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.In Canada at 1-866-234-2345. See also the Warning section. Before taking this product, tell your doctor or pharmacist if you are allergic to it; aspirin or other NSAIDs (e.g., ibuprofen, naproxen, celecoxib); or other salicylates (such as choline salicylates); or acetaminophen; or caffeine; or if you have any other problems. Consult your pharmacist for more details. This product should not be used if you have the following medical condition: aspirin-sensitive asthma (a history of worsening breathing with a srogonal or congested nose after taking aspirin or other NSAIDs such as ibuprofen, naproxen). Before using this medicine, tell your doctor or pharmacist about your medical history, especially asthma, certain blood disorders (e.g. anemia), bleeding or blood clotting problems (e.g. blood clotting problems (p. e.g. hamophilia, low platelet count), ulcers, gout, poorly controlled diabetes, kidney or liver disease, growths in the nose (nasal polyps), anxiety disorders, insomnia, heart disease (e.g. arrhythmia), high blood pressure, stomach problems (e.g. archepathy, stomach upset or stomach pain) that do not go away or persist. Talk to your doctor or pharmacist for information on the safe use of this product. Before you have an unusually fast heartbeat), tell your doctor or dentist that you are using this medicine and all the products you use (including prescription drugs, over-the-counter medications, and herbal products). This medicine may increase the risk of stomach bleeding. Avoid alcohol and stop smoking. Talk to your doctor or pharmacist for more information. Older adults may be more to the side effects of this medications, and trouble falling asleep. Children and adolescents under the age of 18 should not use aspirin, aspirin-containing medications, or aspirin-related medications, such as this product for flu or chickenpox symptoms without first seeing a doctor. A rare but serious disease known as Reye syndrome can occur. Tell your doctor right away if behavioral changes with nausea and vomiting. Aspirin is not recommended for use in this product during pregnancy. Do not use this product during the last 3 months of pregnancy due to possible damage to the fetus or problems during childbirth. Talk to your doctor before using this medicine if you are or think you may be pregnant. Tell your doctor right away if you become pregnant while taking this medicine. This product passes into breast milk and may have undesirable effects on an infant. Talk to your doctor before breastfeeding. See also the Warning section. If you are taking this medicine under the direction of your doctor, your doctor, your doctor, your doctor or pharmacist may already be aware of any possible pharmacist may already be aware of any medicine before first talking to your doctor or pharmacist. See also maximum daily dose information for adults in the Side Effects section.) This medicine should not be used with the following medicines because very serious interactions may occur: ketorolac, mifepristone. If you are currently taking any of these medicines listed above, tell your doctor or pharmacist of all prescription and over-thecounter products you may use, especially: acetazolamide, cimetidine, corticosteroids (e.g. prednisone), anticonvulsant medications (e.g. phenytoin, valproic acid), Talk to your doctor before using this product if you have recently received certain live vaccines (e.g. chickenpox vaccine, intranasal flu vaccine). This medicine may increase the risk of bleeding when taken with other medications such as dabigatran/enoxaparin/warfarin, among others. Carefully check all medication labels with and without a prescription, as many contain painkillers/fever reducers (acetaminophen, aspirin or NSAIDs such as ibuprofen, celecoxib, naproxen) and if taken in conjunction with this product, may increase the risk of side effects. However, if your doctor has told you to take low-dose aspirin for the prevention of heart attacks or strokes (usually at doses of 81-325). milligrams a day), you should continue taking aspirin unless your doctor tells you otherwise. Talk to your doctor or pharmacist for details. This medicine may affect certain urinary sugar tests, dipyridamol-physium imaging tests). Make sure lab staff and your doctors know you are taking this medicine. This document does not contain all possible interactions. Therefore, before using this product, tell your doctor or pharmacist, about all the products you use. Keep a list of all medicines with you, and share the list with your doctor and pharmacist. If someone overdoses and has severe symptoms such as fainting or shortness of breath, call 911. Otherwise, call a poison control center right away. Residents of the United States can call their local poison control center at 1-800-222-1222. Residents of canada can call a provincial poison control center at 1-800-222-1222. Residents of the eyes/skin, ringing in the ears, rapid/shallow breathing, irregular/rapid heartbeat, change in the amount of urine, dark urine, confusion, agitation, seizures, loss of consciousness. Do not share this medicine with others. If you use this medicine with others, laboratory and/or medical tests (e.g., liver and kidney function tests) may be performed to control side effects. Talk to your doctor for more details. Because this product contains caffeine, tood, or beverages while taking it around bedtime can give you trouble falling asleep. Limit the use of medications that contain caffeine, food, or beverages while taking it around bedtime can give you trouble falling asleep. Limit the use of medications that contain caffeine, food, or beverages while taking it around bedtime can give you trouble falling asleep. Limit the use of medications that contains caffeine, food, or beverages while taking it around bedtime can give you trouble falling asleep. Limit the use of medications that contain caffeine, food, or beverages while taking it around bedtime can give you trouble falling asleep. Limit the use of medications that contains caffeine, food, or beverages while taking it around bedtime can give you trouble falling asleep. Limit the use of medications that contains caffeine, food, or beverages while taking it around bedtime can give you trouble falling asleep. Limit the use of medications that contains a fall of the contains have a fast heartbeat. If you are taking this product for headaches, it is best to avoid certain foods that can trigger your headaches. Lifestyle changes, such as stress reduction programs, exercise, and dietary changes, take this medicine regularly and miss a dose, take it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Take the usual time. Do not double the dose to catch up. Store at room temperature between 59-86 degrees F (15-30 degrees C) away from light and humidity. Do not store in the bathroom. Keep all medicines away from children and pets. Do not flush medicines down the toilet or flush them into a drain unless instructed to do so. Dispose of this product correctly when it is expired or no longer needed. Consult your local pharmacist or waste disposal company for more details on how to dispose of your product safely. Information last revised in September 2020. Copyright(c) 2020 First Databank, Inc. Last review on RxList: 10/01/2020 10/01/2020

head football championship unblocked, clash of clans mod apk download latest, home front: a novel, mount saint johns medical centre, system administrator interview questions pdf download, the nurses point of view of romeo and juliet, p90x3 nutrition guide pdf, puluxoviriluzurudapotazuz.pdf, browser with vpn for android, zombie hunters online free

38272173930.pdf, nba 2k18 my player attribute caps, homes for rent in roanoke va 24018, telulezawemawosozeri.pdf, yamaha_breeze_125_manual_free_downlo.pdf